

TOLERANCE OF BARLEY (*HORDIUM VULGARE* L.) TO CHLORIDES OF SODIUM, CALCIUM AND MAGNESIUM

I. Rajpar¹, M. Sultan¹, Zia-ul-hassan¹, N. Debar² and M. A. Kharal³

¹Centre for Biosaline Agriculture (CBSA), Sindh Agriculture University,
Tandojam, Pakistan

²Soil Science Division, Nuclear Institute of Agriculture, Tandojam, Pakistan

³Department of Soil Science, Bahauddin Zakarya University Multan, Pakistan

ABSTRACT

The effect of five chloride salt compositions on the growth and yield of *Hordium vulgare* L. (cv. local) was compared in a pot experiment designed with three replications. Five chloride salt compositions included in the study were: T₁: Control (without salt); T₂: NaCl; T₃: MgCl₂; T₄: CaCl₂ and T₅: Equal concentration (40+40+40 mM) of NaCl: MgCl₂: CaCl₂. The concentration of each salt composition was 120 mM. The results obtained from the study indicated that the influence of 120 mM NaCl was more lethal to barley plants. Compared to other salt compositions, plants stressed with NaCl had higher concentration of Na⁺ and lower concentration of K⁺ both in straw and grains, which exhibited lower K⁺/Na⁺ ratio.

Keywords: Barley, calcium, chlorides, sodium, magnesium.

INTRODUCTION

Soil salinity decreases plant growth by imposing hyper-osmotic and hyper-ionic effects (Jaleel *et al.*, 2007). Among the specific toxic ions, Cl⁻ has been found as one of the most dominant toxic anions present in the saline soils of Sindh (Sial, 1985; Rajpar and Sial, 1997). Chloride particularly disturbs the physiology of plants especially by reducing the uptake of nutrient anions such as nitrates, phosphates, etc. (Munns and Tester, 2008). Availability of excessive chloride concentration in root medium induces diverse effects on crops yield, net photosynthesis rate (Boyer, 1965; Rajasekaran *et al.*, 1997), chlorophyll content, respiration, transpiration, pre-dawn leaf water potential, stomatal density and conductance, leaf area (Katerji *et al.*, 2000; Romero-Aranda *et al.*, 2001), CO₂ fixation, protein synthesis (Ramagopal, 1987; Levine *et al.*, 1990) and other essential biochemical and metabolic processes of the plant.

Corresponding author: irajpar@yahoo.com

Presence of various forms of chlorides such as NaCl, CaCl₂, MgCl₂, etc. may differ in their nature to affect growth and development of plants. Chlorides of Na⁺ may become more toxic and injurious to plants than the Cl⁻ of Ca²⁺ and Mg²⁺. At an equal concentration, the mixture of chlorides of Na⁺, Ca²⁺ and Mg²⁺ can be less toxic and injurious to plants than the Cl⁻ of Na⁺ alone. There are several reports which indicate that the plants differ in their tolerance to salinity (Kumar *et al.*, 2002) suggesting that plants probably differ in tolerance to chloride toxicity. Some crop species and varieties can be found much more tolerant or sensitive to chlorides than the others. Also the higher sensitivity of cereals to salinity has been better explained on the basis of Cl⁻ accumulation rather than Na⁺ (Qadir, 1990). It is therefore essential to identify important commercial crops including cereals tolerant to chloride stress rather than Na⁺. Among the cereals, barley is considered as one of the important grains of Pakistan used for forage, grain and industrial purpose, including distilled beverages. Although, considerable research on salt-tolerance of barley has already been done (Naseer *et al.*, 2001; Ahmed *et al.*, 2003; Bagci *et al.*, 2003); but to the knowledge of authors very little attention has so far been paid to the tolerance of barley plants to various chloride salt compositions. This study reports the tolerance of barley plants against various chloride salt compositions.

MATERIALS AND METHODS

This experiment was conducted in an open environment (wire-house) using black colored plastic pots with capacity of 10 kg soil/pot and drainage holes at the bottom. The pots were placed on wooden benches of the wire-house and arranged in a CRD design. The soil filled in the pots was collected from the field previously cropped to cotton. The soil test results (CBSA Research Laboratory) indicated that it was clay loam in texture (26.30% sand, 35.09% silt and 38.61% clay), non-saline (ECe 1.04 dS m⁻¹) and non-sodic (ESP 4.89) in nature, slightly alkaline (pH 7.6), moderately calcareous (CaCO₃ 9.32%) in reaction to HCl, low in organic matter (0.97%) and total nitrogen (0.049%) contents. However, it was rich in extractable potassium (178.28 mg kg⁻¹). After overnight soaking barley seeds of local variety, obtained from Wheat Section of Agriculture Research Institute (ARI), Tandojam were sown in the soil. Twenty days after sowing (DAS) seedlings were thinned to five plants per pot and 6 cm apart.

Five chloride salt compositions included in the study were: T₁ = Control (without salt); T₂ = 120 mM NaCl T₃ = 120 mM MgCl₂; T₄ = 120 mM CaCl₂ and T₅ = 40 mM NaCl + 40 mM CaCl₂ + 40 mM MgCl₂ (mixture). Salt solution for each treatment was prepared by dissolving the required salt in the distilled water, following the method of Rowell (1994). In the control treatment, plants were only receiving fresh water, whereas in the salt treatments, plants were established with salt solutions. The nitrogen was applied @160 kg N ha⁻¹ through urea and diammonium phosphate (DAP), phosphorus was applied @ 80 Kg P₂O₅ ha⁻¹ through DAP and potassium was applied @ 80 kg K₂O ha⁻¹ through sulfate of potash (SOP). Full dose of phosphorus and potash was applied at the time of sowing, nitrogen was applied in three splits i.e. 1/2 at sowing, 1/3 at 1st irrigation and 1/3 at booting stage.

At physiological maturity plants from each treatment were harvested by cutting with sharp sickle at soil level. The data on plant height (cm) and number of tillers per plant, were recorded. The ear-heads were separated from the straw. The grains were separated by threshing with hands. The data on grain and straw yields per plant were recorded. The straw and grains were then used to determine Na⁺ and K⁺ contents and calculate the ratio of Na⁺ to K⁺. The plant data were processed for the analysis of variance (ANOVA) using MINITAB statistical software version-13. Treatment means were compared using SED and LSD tests at 5% probability level.

RESULTS AND DISCUSSION

The results presented in the Table 1 indicate the effect of various chloride salt composition on plant height, number of tillers per plant and straw and grain yields per plant. It is evident from the data that compared to control, the plants stressed with NaCl, MgCl₂, CaCl₂ and equal mixture of these salts were about 10, 7, 7 and 9% shorter in height, developed 30, 25, 24 and 28 % fewer tillers plant⁻¹, showed 49, 43, 33 and 47% reduction in grain yield per plant and 51, 42, 39 and 48 % reduction in straw yield per pant. It appears that compared to Ca²⁺ and Mg²⁺ ions, the Cl⁻ are more toxic to barley plants in presence of Na⁺ ions. As the plants stressed with NaCl and mixture of salts having only 40 mM Na⁺ showed much more reductions in almost all growth and yield traits. That was possibly the phytotoxicity jointly created by both Na⁺ and Cl⁻ ions for barley plants in the treatments where NaCl was added. In addition to that plants were also facing dual problem of salinity i.e osmotic effect due to 120 mM concentration and specific ion effect due to presence of Na⁺ and Cl⁻. There are several evidences which indicate that Ca²⁺ detoxifies the effect of toxic ions including Na⁺, particularly the Ca²⁺ improves the function of cell membrane (Lin, 2014). The results related to ion contents (Tables 2 & 3) obtained from this study also showed the association of toxic ions like Na⁺ and Cl⁻ determined in grain and straw samples with growth and yield data. The plants stressed with NaCl and salts mixture accumulated more Na⁺ and Cl⁻ and less K⁺ both in grain and straw, which also brought down the ratio of K⁺ to Na⁺ ions. This suggests that Ca²⁺ and Mg²⁺ dilute the adverse effect of Na⁺ for plants mainly by improving the physiology of plants.

Table 1. Effect of different chloride salts composition on the growth and yield of barley plants.

Salt composition/ treatments	Plant height (cm)	No. of tillers (plant ⁻¹)	Grain yield (g plant ⁻¹)	Straw wt. (g plant ⁻¹)
T ₁ : Control (without salt)	49.75	4.58	5.76	6.67
T ₂ : NaCl (120 mM)	44.59	3.21	2.96	3.30
T ₃ : MgCl ₂ (120 mM)	46.08	3.33	3.30	4.06
T ₄ : CaCl ₂ (120 mM)	46.19	3.50	3.86	3.50
T ₅ : Mixture of NaCl + CaCl ₂ + MgCl ₂ (120 mM)	45.16	3.31	3.03	3.26
SED	1.98	0.69	0.51	1.82
LSD (0.05%)	4.41	1.53	1.15	4.08

Table 2. Effect of different chloride salts on Na⁺ and K⁺ contents (%) and K⁺/Na⁺ ratio determined in the grains of barley.

Salt composition/ treatments	Na ⁺ (%) in grains	K ⁺ (%) in grains	K ⁺ /Na ⁺ in grains	Ca ²⁺ (%) in grains	Mg ²⁺ (%) in grains	Cl ⁻ (%) in grains
T ₁ : Control (without salts)	0.60	2.12	3.53	0.28	0.29	0.46
T ₂ : NaCl (120 mM)	1.69	0.89	0.53	0.27	0.30	0.84
T ₃ : MgCl ₂ (120 mM)	0.68	1.14	1.67	0.47	0.48	0.59
T ₄ : CaCl ₂ (120 mM)	0.65	1.21	1.86	0.65	0.27	0.61
T ₅ : Mixture of NaCl + CaCl ₂ + MgCl ₂ (120 mM)	0.89	0.92	1.04	0.58	0.31	0.69
SED	0.04	0.04	0.177	0.20	0.037	0.04
LSD (0.05%)	0.10	0.10	0.396	0.045	0.084	0.089

Table 3. Effect of different chloride salt composition on ion content (%) and K⁺/Na⁺ ratio determined in straw of barley.

Salt treatments	Na ⁺ (%) in straw	K ⁺ (%) in straw	K ⁺ /Na ⁺ in straw	Ca ²⁺ (%) in straw	Mg ²⁺ (%) in straw	Cl ⁻ (%) in straw
T ₁ : Control (without salts)	0.61	2.23	3.64	0.24	0.26	0.39
T ₂ : NaCl (120 mM)	2.12	0.92	0.44	0.22	0.27	0.70
T ₃ : MgCl ₂ (120 mM)	0.68	1.20	1.75	0.39	0.44	0.49
T ₄ : CaCl ₂ (120 mM)	0.64	1.28	1.99	0.55	0.27	0.51
T ₅ : Mixture of NaCl + CaCl ₂ + MgCl ₂ (120 mM)	1.12	1.02	0.92	0.48	0.26	0.57
SED	0.073	0.030	0.007	0.015	0.036	0.033
LSD (0.05%)	0.162	0.067	0.016	0.031	0.081	0.074

CONCLUSION

The study presented here demonstrates that Na either alone or in mixtures is much more toxic to rice plants than CaCl₂ and MgCl₂.

REFERENCES

- Ahmed, A. N., U. H. J. Intshar, A. Shamshad and A. Muhammad. 2003. Effect of Na₂SO₄ and NaCl salinity levels on different yield parameters of barley genotypes. *Intl. J. Agric. Biol.*, 5 (2): 157-159.
- Bagci, S. A., H. Ekiz and A. Yilmaz. 2003. Determination of the salt-tolerance of some barley genotypes and characteristics affecting tolerance. *J. Plant Physiol.*, 27: 253-260.
- Boyer, J. S. and E. B. Knipling. 1965. Isopiestic technique for measuring leaf water potentials with a thermocouple psychrometer. *Proc. Natl. Acad. Sci. USA*, 54:1044-1051.

- Jaleel, C. A., P. Manivannan, B. Sankar, A. Kishorekumar and R. Panneerselvam. 2007. Calcium chloride effects on salinity-induced oxidative stress, proline metabolism and indole alkaloid accumulation in *Catharanthus roseus*. C. R. Biologies, 330: 674-683.
- Katerji, N., J. W. Van Hoorn, A. Hamdy and M. Mastrorilli. 2000. Salt-tolerance classification of crops according to soil salinity and to water stress day index. Agri. Water Management, 43: 99-109.
- Levine, R. L., D. Garland, C. Oliver, A. Amici, I. Clement, A. G. Lenz, B. W. Ahn, S. Shaltiel, and E. R. Stadtman. 1990. Determination of carboxyl content in oxidatively modified proteins. Methods Enzymol., 186: 464-478.
- Lin, H., W. Du, Y. Yong, K. S. Schumaker and Y. Guo. 2014. A calcium-independent activation of the Arabidopsis PKS 24, Kinase by its interacting SCA1 Calcium sensor. Plant Physiol. Preview, 1-28 [www. Plantphysiol. Org., Accessed on April 4, 2014].
- Munns, R. and M. Tester. 2008. Mechanisms of salinity tolerance. Annul Review Plant Biology, 59: 651-681.
- Naseer, S., A. Nisar and M. Ashraf. 2001. Effect of salt stress on germination and seedling growth of barley (*Hordeum vulgare* L.). Pak. J. Biol. Sci., 4 (3): 359-360.
- Rajasekaran, L. R., P. E. Kriedemann, D. Aspinall and L. G. Paleg. 1997. Physiological significance of proline and glycinebetaine maintaining photosynthesis during NaCl stress in wheat. Photosynthetica, 34: 357-366.
- Rajpar, I. and N. B. Sial. 1997. Relationship between salts of soil profile and underground water. Pak. J. Agric. Agril. Engg. Vet. Sci., 13: 19-23.
- Ramagopal, S. 1987. Salinity induced tissue specific proteins in barley seedlings. Plant Physiol., 84: 324-331.
- Romero-Aranda, T. Soria and J. Cuartero. 2001. Plant water uptake and plant-water relationships under saline growth conditions. Plant Sci., 160: 265-272.
- Rowell, D. L. 1994. Soil Science Methods and Applications. Addison Wesley Longman Ltd, Edinburgh Gate, Harlow, Essex, Cm2 0J.E, England.
- Sial, N. B. 1985. The formation of salts in the Sind region. Engg. Tech. Res. J., 4: 24-28.

(Received November 19, 2013; Accepted February 28, 2014)